

**WARNING**  
Keep out of reach of children

# Turbostar 430SC

**FUNGICIDE**

**ACTIVE INGREDIENT:**

Contains 430 g/L Tebuconazole in the form of a suspension concentrate.

**A systemic triazole (DMI) fungicide for the control of diseases in cereals, grass seed crops, peas and onions, for the control of brown rot in stonefruit and to control rust and fusarium in pasture.**

Registered pursuant to the ACVM Act 1997, No. P9962

See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

Approved pursuant to the HSNO Act 1996, Approval Code HSR007675.

See [www.epa.govt.nz](http://www.epa.govt.nz) for approval controls.

**IN A TRANSPORT EMERGENCY: CALL 111, POLICE OR FIRE BRIGADE**

**AgSta**  <sup>®</sup>



**READ THE LABEL COMPLETELY BEFORE MIXING OR APPLYING  
DIRECTIONS FOR USE:**

**It is an offence to use this product on animals.**

For field crops ensure complete coverage by applying at least 200 litres water/ha by ground, or 50 litres water/ha by air. Apply preventatively when disease levels are low but disease risk is high.

For stonefruit the rates suggested are for high volume spraying to run off. For concentrate spraying adjust dilution rate accordingly.

Rainfall within one hour following application may reduce effectiveness. If 15mm or more rain falls within 3 hours of treatment, pasture should be re-sprayed.

<b>CROP</b>	<b>DISEASE</b>	<b>RATE</b>	<b>NOTES</b>
Barley	Ear disease complex, Leaf rust, Net blotch (spot and net form), Ramularia, Scald, Spot blotch	440mL/ha	Apply at first appearance of disease and repeat 3-4 weeks later if disease pressure persists. For the ear disease complex, apply to fully emerged ears. Apply a maximum of three DMI applications, including seed treatment, per crop. Where powdery mildew is expected to be severe, use DMI's in conjunction with alternative fungicides.
Wheat	Glume blotch, Leaf rust, Powdery mildew, Speckled leaf blotch, Stripe rust, Ear disease complex		
Oats	Crown rust		
Onions	White rot	875mL/ha	Apply in 1,000 litres water/ha. Make the first application 6-8 weeks after sowing and follow with two further applications at 3-4 week intervals.
Peas	Powdery mildew	145mL/ha	Apply preventatively at flowering or at first sign of disease. Repeat if necessary 14 days later.
Ryegrass Seed Crops	Crown rust, Stem rust	440mL/ha	Apply at first appearance of disease and repeat 3-4 weeks later if necessary.

	Blind seed disease	440mL/ha	Apply at the start of flowering and repeat 10-14 days later.
Stonefruit	Brown rot (Blossom blight)	23mL/100L water	(a) Apply at 7-10 day intervals from pink to shuck fall. Shorter intervals may be needed in wet weather. Apply non-DMI fungicides in the preharvest period. or (b) Apply 2-3 applications at 7-10 day intervals in the preharvest period. Use non-DMI fungicides earlier in the season. Do not apply more than 3 applications per season, preferably in mixture with a non- DMI fungicide.
Pasture (grass/clover)	Rust Fusarium	440mL/ha	Apply at first sign of disease. Ensure complete coverage of grazed areas. A second application may be necessary if conditions favour disease development. Do NOT reapply to the same area at less than a 28 day interval, or more than twice per season. Do NOT apply when temperatures are above 20°C or wind speed greater than 12kph. <b>Ground:</b> Apply in 100 to 200 litres of water/ha. Add organosilicone at 100mL/100 L <b>Aerial:</b> Apply in 50 litres water/ha. Add organosilicone at 200mL/100L.

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

## **Withholding Periods:**

Cereals for grain – 49 days

Fodder cereals – 28 days

Onions – 35 days

Peas – 14 days

Ryegrass seed crops – 30 days

Stonefruit – 1 day. For export crops check with Summerfruit NZ or your exporter.

Pasture – Animals (including lactating animals) may be reintroduced into treated pasture 10 days after application. Animals for slaughter must be placed on clean pasture for 6 days prior to slaughter. Pasture is considered clean from residues 45 days after treatment. Where pasture cannot be guaranteed clean, such as in aerial applications, animals must not be sent for slaughter for 51 days after treatment of the pasture.

## **Resistance Management**

<b>GROUP</b>	<b>03</b>	<b>FUNGICIDE</b>
--------------	-----------	------------------

**Turbostar 430SC** contains a fungicide from the triazole group of fungicides and has the DMI (Demethylation Inhibitors) mode of action. Resistance to this fungicide could develop from excessive use. To minimise this risk use strictly in accordance with label instructions and resistance management strategies that exist for any fungus listed on the label. Refer to [www.resistance.nzpps.org](http://www.resistance.nzpps.org) for further resistance management advice.

## **CONDITIONS OF SUPPLY**

**READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING.**

**IF THESE CONDITIONS ARE UNACCEPTABLE, RETURN AT ONCE UNOPENED.**

This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer or registrants control. The manufacturer and registrant decline all liability for damage resulting from improper storage or use of the product.



**Registrant: AGROBEATS TECH CO., LTD**

C-47, No 69, Shuangfeng Road, Pukou Economic Development Zone,  
Pukou District, Nanjing, P R China 210031

**NZ Agent: AgStar New Zealand Pty Ltd**

4 Putakitaki Street, Lincoln, 7608

[www.agstar.co.nz](http://www.agstar.co.nz)

**Distributed by: Farmlands Co-operative Society Limited**

535 Wairakei Road, Burnside, Christchurch 8053

TEL: 0800 200 600

©AgStar is a Registered Trademark of Agstar New Zealand Pty Ltd