

WARNING
KEEP OUT OF REACH OF CHILDREN
ECOTOXIC

Forestar Solo

HERBICIDE

ACTIVE INGREDIENT:

500g/L terbuthylazine in the form of a suspension concentrate

**For selective weed control in forestry and non-selective residual weed control in non-cropland areas
Not to be applied to pasture or cropland that may be grazed by food-producing animals**

Approved pursuant to the HSNO Act 1996, Approval Code HSR100978
See www.epa.govt.nz for approval controls.

IN A TRANSPORT EMERGENCY: CALL 111, POLICE OR FIRE BRIGADE



READ LABEL COMPLETELY BEFORE OPENING AND USING PRODUCT SHAKE OR STIR WELL BEFORE USE

GENERAL INSTRUCTIONS

Forestar Solo is a selective herbicide for use in forestry, and a non-selective herbicide for residual weed control in noncropland areas. This product is absorbed by the roots, although some foliar absorption does take place. Do not use on pasture or cropland.

MIXING

Stir or mix well. Add the full quantity of product to the partly filled spray tank while agitating. Fill tank and agitate to ensure thorough mixing and continue agitation while spraying. Agitate vigorously from the bottom if allowed to stand

APPLICATION

Forestar Solo may be applied through all conventional spray equipment. Do not use strainers finer than 300 microns.

COMPATIBILITY

This product is compatible with most tank additives including amitrole, and glyphosate. It is advisable to add **Forestar Solo** to the partly filled spray tank first, mix and then incorporate the additive with agitation.

RAINFALL

Rainfall within 3 hours following treatment may reduce effectiveness. Earlier re-treatment should be considered following heavy rain.

DIRECTIONS FOR USE

Forestar Solo controls a wide range of weeds including barley grass, brown top, catsear, chickweed, clovers, cocksfoot, fathen, Kentucky bluegrass, nodding thistle, poa species, redroot, ryegrass species, Scotch thistle, seedling gorse, Shepherd's purse, storksbill, sweet vernal, willow weed, Yorkshire fog plus annual grasses and broadleaf weeds. The product does not control deep rooted perennial grasses such as paspalum, kikuyu or Indian doab. It is recommended to add Glyphosate 510SL to the mix at the label rate to control these weeds.

APPLICATION RATE

The maximum application rate is 9.9kg of terbuthylazine/ha.

Crop	Application method	RATE	Critical Comments
Radiata Pine Douglas fir	Aerial	15-19L/ha in 150-300L water	Apply prior to spring growth. Use the higher rate on perennial grasses or when vegetation is dense.
	Kanpsack application	1.5-1.9mL/square metre	Apply with sufficient water to obtain good coverage.
Non-Cropland Areas	Broadcast Application	15-19L/ha in 150-300L water	Use sufficient water to obtain even coverage. Add AGPRO Glyphosate to control deep rooted perennial weeds.

RESISTANT WEEDS WARNING

GROUP	5	HERBICIDE
-------	----------	-----------

Forestar Solo contains a triazine herbicide that inhibits photosynthesis at photosystem II. Naturally occurring resistant biotypes are known to exist and can become more prevalent after many years of continuous or excessive use of this product. To prevent or delay resistance use **Forestar Solo** in tank mixes as appropriate, and/or in rotation with herbicides that have a different mode of action

CONDITIONS OF SALE

READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING.

IF THESE CONDITIONS ARE UNACCEPTABLE, RETURN AT ONCE UNOPENED.

This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer or registrants' control. The manufacturer and registrant decline all liability for damage resulting from improper storage or use of the product.



Registrant: AgStar New Zealand Pty Ltd

4 Putakitaki Street, Lincoln, 7608

Distributed by: Farmlands Co-operative Society Limited

535 Wairakei Road, Burnside, Christchurch 8053

TEL: 0800 200 600

©AgStar is a Registered Trademark of Agstar New Zealand Pty Ltd