

DANGER

KEEP OUT OF REACH OF CHILDREN

ECOTOXIC

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Access 840

HERBICIDE

ACTIVE INGREDIENT:

840 G/LITRE ACETOCHLOR IN THE FORM OF AN EMULSIFIABLE CONCENTRATE. ALSO CONTAINS 108 G/L OF 2,2-DICHLORO-N,N-DI-2-PROPENYLACETAMIDE AS A HERBICIDE SAFENER.

GROUP	15	HERBICIDE
--------------	-----------	------------------

FOR THE SELECTIVE PRE-EMERGENCE CONTROL OF ANNUAL GRASSES AND MANY BROADLEAF WEEDS IN GRAIN AND SILAGE MAIZE AND SWEETCORN.

Registered pursuant to the ACVM Act 1997, No. P10214

Approved pursuant to the HSNO Act 1996, Approval Code HSR000404.

IN A TRANSPORT EMERGENCY: Dial 111 for POLICE or FIRE

IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING THIS PRODUCT

AgSta  [®]

DIRECTIONS FOR USE

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

GRAIN AND SILAGE MAIZE AND SWEETCORN: Rate and Timing of Application: 2.5 – 3.0 litres in at least 200 litres of water per hectare to a freshly prepared seedbed immediately after planting and before crop emergence. Use the higher rate in clay soils and those with higher organic matter. If desired, Atrazine may be added to the tank mix to improve the spectrum of activity. Alternatively, Atrazine or Dicamba may be applied as follow-up post emergence treatment.

MIXING: To ensure mixing pour the correct amount into the partly filled tank and add the remainder of the water while agitating. Agitate thoroughly before and during spraying and before spraying after a stoppage.

COMPATIBILITY: Compatible with Atrazine.

SOIL CONDITIONS/RAINFALL: Best results will be obtained when application is made to a freshly prepared, fine, moist soil, free of trash, with at least 10 mm of rain within 1 week of application. Under drier soil conditions, 30 mm of rain within 2 – 3 weeks is necessary. Application by pre-plant incorporation may be beneficial under very dry conditions.

SOIL RESIDUES: Under normal use, soil residues should not carry over to successive crops. In a replant situation when the previous treated crop may have failed for whatever reason, get technical advice before

replanting. If establishing pasture in peat soils, cultivate soil before sowing if **Access 840** has been applied to the previous crop.

RE-ENTRY PERIOD: When fully dry.

RESISTANCE MANAGEMENT

GROUP	15	HERBICIDE
-------	----	-----------

Access 840 contains acetochlor, a Group 15 herbicide belonging to the alpha-chloroacetamide chemical family. Resistance to this and other chloroacetamide or oxyacetamide herbicides could develop from excessive use. To minimise this risk, use strictly in accordance with label instructions and rotate use with herbicides with a different mode of action. Consult <http://resistance.nzpps.org/index.php> for alternative modes of action and details of resistance management strategies (if any) for the crops listed on this label.

CONDITIONS OF SUPPLY:

READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING. IF THESE CONDITIONS ARE UNACCEPTABLE, RETURN AT ONCE UNOPENED.

To the extent permitted by law Agrobeats does not accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with these goods, including without limitation their storage, application, handling or use. Agrobeats does not make any warranties (express or implied) with respect to fitness for purpose, merchantable quality or any other matter. To the extent

only that Agrobeats is prevented by law from contracting out of the Consumer Guarantees Act, then these conditions apply subject to that Act. The Act shall not apply where these goods are acquired for business purposes.



AgStar New Zealand Pty Ltd

4 Putakitaki Street, Lincoln, 7608

Tel 027 483 9756

®AgStar is a Registered Trademark of Agstar New Zealand Pty Ltd